

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 August 2002 (08.08.2002)

PCT

(10) International Publication Number
WO 02/061067 A1

(51) International Patent Classification⁷: C12N 9/96, 9/20

(21) International Application Number: PCT/NL02/00079

(22) International Filing Date: 1 February 2002 (01.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1017258 1 February 2001 (01.02.2001) NL

(71) Applicant (*for all designated States except US*): **TECHNISCHE UNIVERSITEIT DELFT** [NL/NL]; Julianalaan 134, NL-2628 BL Delft (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LÓPEZ SER-RANO, Paloma** [ES/NL]; Roodborstlaan 37, NL-2623 HA Delft (NL). **CAO, Linqiu** [CN/NL]; Aart van der Leeuwlaan 197, NL-2624 PP Delft (NL). **VAN RANTWIJK, Frederik** [NL/NL]; Groensvoorde 104, NL-2742 DL Waddinxveen (NL). **SHELDON, Roger, Arthur** [GB/NL]; Regentesselaan 13, NL-2281 VA Rijswijk (NL).

(74) Agent: **ALTENBURG, Bernardus, Stephanus, Franciscus**; Octrooibureau Los En Stigter B.V., Weteringschans 96, NL-1017 XS Amsterdam (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A METHOD OF PREPARING CROSS-LINKED ENZYME AGGREGATES

(57) Abstract: The invention relates to a method of preparing cross-linked enzyme aggregates (CLEAs) comprising the steps of a) precipitating a dissolved enzyme present in a liquid medium; b) cross-linking with a cross-linker, and c) optionally, washing the cross-linked enzyme aggregate. According to the invention at least one treatment is performed chosen from the group consisting of i) performing step a) in the presence of a compound selected from the group consisting of i) a crown ether, and ii) a surfactant, and II) performing step c) with a second liquid differing in organic composition from the liquid.



WO 02/061067 A1

A method of preparing cross-linked enzyme aggregates

The present invention relates to a method of preparing cross-linked enzyme aggregates comprising the steps of
a) precipitating a dissolved enzyme present in a liquid medium;

- 5 b) cross-linking with a cross-linker, and
c) optionally, washing the cross-linked enzyme aggregate

Such a method is generally known in the art. It is used to prepare enzyme preparations for performing various enzyme-catalysed reactions. For example, Cao L. et al. (Organic Letters vol. 2 (10) p. 1361-1364, 2000) describe the
10 preparation of penicillin acylase-based aggregates. The cross-linked aggregates disclosed are less active than the native enzyme.

The object of the present invention is to provide a
15 method making it possible to achieve aggregates with increased enzymatic activity.

To this end, the invention is characterized in that at least one treatment is performed chosen from the group consisting of I) performing step a) in the presence of a compound selected from the group consisting of i) a crown ether,
20 and ii) a surfactant, and II) performing step c) with a second liquid differing in organic composition from the liquid.

Applicants have surprisingly found that such a treatment may result in an cross-linked enzyme aggregate
25 (CLEA) with an enhanced activity. While the present invention does neither guarantee this enhanced activity for a particular enzyme used to catalyse a particular reaction, nor allows for a prediction of an enhanced activity, it does provide for an extra parameter to be controlled to achieve a better aggregate. That is, for a particular enzyme, it may be investigated with little effort and using simple and routine experimentation whether its catalytic activity can be enhanced using the treatment. From a range of enzymes capable of catalyzing the desired reaction, the best CLEA can easily and
35 with little effort be selected. Because the CLEA is insoluble, it can be washed with a second liquid which is com-

pletely aqueous, organic, or a mixture of the two. After the treatment, the CLEA may be lyophilized or stored as a suspension for future use.

Overbeeke P.L.A. et al. Describe in J. of Mol. Cat.
5 B: Enzymatic 10, p. 385-393 (2000) a method in which a lipase is crystallized under the influence of PEG in the presence of 2-methyl-2,4-pentanediol, and crystals formed are cross-linked. In contrast, the present invention makes use of a precipitating agent to form aggregates instead of crystals.

10 Preferably, step a) and step b) are performed simultaneously.

This allows for a simplified procedure, requiring less time and effort.

According to a preferred embodiment, the compound
15 is removed after cross-linking.

It is not necessary that the surfactant and or crown ether are present during the catalytic reaction to be performed by the CLEA to maintain the high activity. Thus the compound may be removed, in case it may interfere with the
20 reaction to be catalysed, or for other reasons. Removal may be achieved using any method known in the art, and in particular through dialysis or washing using centrifugation.

According to an important application of the method according to the invention, the enzyme is a lipase.

25 Hence, the invention also relates to a method for performing a lipase-catalysed reaction, said reaction being chosen from the group consisting of i) transesterification; ii) interesterification; iii) hydrolysis; iv) esterification; v) ammoniolysis; vi) aminolysis; and vii) perhydrolysis,
30 which method is characterized, in that a cross-linked lipase-aggregate is prepared according to the invention, and said cross-linked lipase-aggregate is contacted with a substrate.

The present invention will now be illustrated with reference to the following examples.

35

EXAMPLE 1.

Preparation of cross-linked lipases

- *R. miehei* lipase Lipozyme (Novo Nordisk)

- *Candida antarctica* lipase B SP525 (Novo Nordisk)
- *P. alcaligenes* lipase (Gist-brocades).

METHOD Ia) In the presence of (NH₄)₂SO₄ (Water)

0.5 ml stock enzyme solution (Lipozyme or SP525, Novo Nordisk, Copenhagen, Denmark) or 50 mg PaL enzyme powder (PaL is *P. alcaligenes* lipase (Gist-Brocades, Delft, the Netherlands)), are dissolved in 1 ml potassium phosphate buffer (100 mM, pH 7) in a 10 ml centrifuge tube. 550 mg (NH₄)₂SO₄ was added followed by 1 ml of a precipitating solution A consisting of 550 mg (NH₄)₂SO₄ (the precipitant) in potassium phosphate buffer (100 mM, pH 7), and 80 µl glutardialdehyde (25% solution in water) are added. The mixture is left stirring at 4°C (room temperature for SP525) for 17 hours. 3 ml H₂O is added to the mixture and then centrifuged. The supernatant is decanted and the residue is washed, centrifuged and decanted 3 more times with H₂O (5 ml each time). The final enzyme preparation is kept in 5 ml H₂O. If necessary the solid is dispersed by stirring with magnets.

METHOD Ib and Ic) In the presence of (NH₄)₂SO₄ and surfactant

The method Ia) was repeated, with the difference that the precipitating solution A contained in addition 25 mg sodium dodecylsulphate (SDS) (method Ib) or 25 mg Triton X-100 (TR) (method Ic).

METHOD Id) In the presence of dimethoxyethane (DME)

0.5 ml stock enzyme solution (Lipozyme or SP525) or 50 mg PaL enzyme powder, are introduced in a 10 ml centrifuge tube, after which 1 ml potassium phosphate buffer (100 mM, pH 7), 3 ml DME and 80 µl glutardialdehyde are added. The mixture is left stirring at 4°C (room temperature for SP525) for 17 hours.

1 ml DME is added to the mixture and then centrifuged. The supernatant is decanted and the residue is washed, centrifuged and decanted 3 more times with DME (5 ml each time). The final enzyme preparation is kept in 5 ml DME. If necessary the solid is dispersed by stirring with magnets.

METHOD Ie) In the presence of DME and a Crownether

Method Id) was repeated, except that an amount of 6.9 mg the crownether dibenzo-18-crown-6 was added too together with the

3 ml DME.

Cross-linked lipase preparations were kept in suspension, either in water or DME, named preparations Ia-e.

5 EXAMPLE 2

Activity tests

The activity of CLEA's prepared according to the preparations Ia-e) were tested with three reactions. These reactions were performed either in an aqueous solution, or in an organic solution.

For tests IIb and IIc, the following general procedure was followed, before performing the assay:

- Of aqueous suspensions, 0.5 ml samples of CLEA suspension were lyophilized in 2 ml Eppendorf cup each.

- 15 - Of organic suspensions, 0.5 ml samples of CLEA suspension were centrifuged in a 2 ml Eppendorf cup each, and the supernatant was decanted.

- Native enzyme, used as a control, was used directly from the stock solution (50 μ l), or as powder (5 mg), without further treatment.

For text IIb and IIc, analyses were performed using a GC Varian Star 3600 gas chromatograph (Varian). Column: CP WAX 52 CB 50 m x 0.53 mm. Carrier gas: nitrogen.

25 *Test IIa) Hydrolysis of p-nitrophenyl propionate (performed in water)*

The enzyme preparation (50 μ l) was suspended in 0.5 ml water (1.5 ml in case of SP525) and 200 μ l (50 μ l in case of SP525) of the suspension was added to 2.5 ml ester solution (0.4 mM p-nitrophenyl propionate in water at 25°C). Measurement of the increase of absorbance at 348 nm (appearance of p-nitrophenol) in the hydrolysis of p-nitrophenyl propionate was monitored using a Cary 3-Bio UV spectrophotometer from Varian. Values are given as percentage of the activity of the native enzyme.

35 The result is summarized in table 1

Table 1

Hydrolysis of p-nitrophenyl propionate in water

Enz	Ia	Ib	Ic	Id	Ie	Native
Lipozyme	152	218	44	0	0	100
SP525	93	40	129	151	177	100

5 % of activity of native enzyme

Test IIb) Transesterification of ethyl octanoate with octanol (performed in diisopropylether)

10 0.5 ml diisopropyl ether is added to the Eppendorf cup containing the enzyme. 50 mg activated zeolite NaA (Aldrich) is added. 0.5 ml solution of 86 mM ethyl octanoate, 400 mM octanol, and 50 mM decane in diisopropyl ether is added and left stirring at room temperature. At the specified time, the re-

15 action mixture is centrifuged, an aliquot (100 μ l) withdrawn, dissolved in 0.5 ml DME, and analyzed by GC.

The result is summarized in table 2

Table 2

20

Transesterification of ethyl octanoate with octanol in i-Pr₂O

Enz	Ia	Id	Ic	Id	Ie	Native
Lipozyme	0.16	3.48	0.33	23.79	18.89	0.16
Pal	0.15	7.11	24.18	0.11	0.27	17.55

Activity expressed as μ mol of octyl octanoate produced in 24 h.

25

Test IIc) Hydrolysis of ethyl octanoate (performed in DME/water mixture)

50 μ l H₂O and 0.95 ml solution of 43 mM ethyl octanoate, and 25 mM decane is added to Eppendorf cup containing the enzyme

30 and left stirring at room temperature. At the specified time, anhydrous Na₂SO₄ is added to the reaction mixture, shaken, and centrifuged. An aliquot (100 μ l) is then withdrawn, dissolved

in 0.5 ml DME, and analyzed by GC.
The result is summarized in table 3

Table 3

5

Hydrolysis of ethyl octanoate in DME/water

Enz	Ia	Ib	Ic	Id	Ie	Native
Lipo- zyme 24h	24.34	<u>40.48</u>	25.37	22.85	22.73	16.34
Pal. 3h	35.82	32.02	<u>40.26</u>	0.49	0.37	3.34

Activity expressed as μmol of octanoic acid produced in 24 or 3 h.

10

As shown by the above results, adding a surfactant or a crown ether in accordance with the present invention, may have a beneficial effect on the enzymatic activity.

15 EXAMPLE 3

Washing test

Methode Ia was repeated using SP525. The supernatant is decanted and the pellet resuspended in 5 ml water, centrifuged and the supernatant is decanted.

20 The pellet obtained is subjected to the same washing procedure with 3 * 5 ml water or DME. After the last washing step, the CLEA is kept in 5 ml of the same solvent.

The two CLEAs were tested using the transesterification of ethyl octanoate with octanol (as described in test IIc).

25 0.5 ml DME is added to 50 μl enzyme preparation in a 2 ml Eppendorf. The enzyme washed with water only was previously lyophilized in the Eppendorf cup. 50 mg activated zeolite NaA is added. 0.5 ml solution of 86 mM ethyl octanoate, 400 mM octanol, and 50 mM decane in DME is added and left stirring
30 at room temperature. After 1 hour, the reaction mixture is centrifuged, an aliquot (100 μl) withdrawn, dissolved in 0.5 ml DME, and analyzed by GC.

Table 4

Transesterification of ethyl octanoate with octanol (in DME)

5

Processing solvent	Activity
Water	5.41
DME	14.03

Activity expressed as μmol of octyl octanoate produced in 1 h.

10 EXAMPLE 4

Lipase CLEAs were prepared according the procedure described in example 1 and were tested in the hydrolysis of methyl mandelate. The enzyme was added to a 50 mM solution of methyl mandelate in dimethoxyethane-potassium phosphate
15 buffer pH 7 (95:5, v/v). After 48 hours the conversion and enantiomeric excess of the substrate (ee_s) were analyzed using HPLC (see Table 5).

Table 5

20

Precipitant	Additive	Conv. (%)	ee_s	E
Dimethoxyethane	-	31.0	0.31 (R)	7
Dimethoxyethane	CR	33.5	0.33 (R)	7
Acetone	-	26.7	0.26 (R)	8
$(\text{NH}_4)_2\text{SO}_4$	SDS	27.9	0.27 (R)	8
$(\text{NH}_4)_2\text{SO}_4$	TR	23.2	0.24 (R)	10
Free enzyme	-	33.9	0.33 (R)	6

CR : Crownether dibenzo-18-crown-6

SDS: Sodium dodecylsulfate

TR : Triton X-100

ee_s : enantiomeric excess of the substrate

25 E : Enantiomeric ratio.

From table 5 it can be seen that the Enantiomeric ratio can be increased using the method according to the present invention.

CLAIMS

1. Method of preparing cross-linked enzyme aggregates comprising the steps of
- 5 a) precipitating a dissolved enzyme present in a liquid medium;
- b) cross-linking with a cross-linker, and
- c) optionally, washing the cross-linked enzyme aggregate, **characterized**, in that at least one treatment is performed chosen from the group consisting of I) performing step a) in
- 10 the presence of a compound selected from the group consisting of i) a crown ether, and ii) a surfactant, and II) performing step c) with a second liquid differing in organic composition from the liquid.
2. Method according to claim 1, **characterized**, in
- 15 that step a) and step b) are performed simultaneously.
3. Method according to claim 1 or 2, **characterized**, in that the compound is removed after cross-linking.
4. Method according to any of the preceding claims, **characterized**, in that the enzyme is a lipase.
- 20 5. Method for performing a lipase-catalysed reaction, said reaction being chosen from the group consisting of i) transesterification; ii) hydrolysis; and iii) ammonioly-sis, **characterized**, in that a cross-linked lipase-aggregate is prepared according to claim 4, and said cross-linked li-
- 25 pase-aggregate is contacted with a substrate.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 02/00079

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N9/96 C12N9/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, MEDLINE, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CAO L ET AL: "Cross-linked aggregates of penicillin acylase: Robust catalysts for the synthesis of beta-lactam antibiotics." JOURNAL OF MOLECULAR CATALYSIS B ENZYMATIC, vol. 11, no. 4-6, 2001, pages 665-670, XP001079507	1
Y	ISSN: 1381-1177 the whole document --- -/-	1-5



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

18 June 2002

Date of mailing of the international search report

15/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Dumont, E

INTERNATIONAL SEARCH REPORT

Int. 1st Application No

PCT/NL 02/00079

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CAO L ET AL: "Cross-linked enzyme aggregates: a simple and effective method for the immobilization of penicillin acylase." ORGANIC LETTERS. UNITED STATES 18 MAY 2000, vol. 2, no. 10, 18 May 2000 (2000-05-18), pages 1361-1364, XP002202388 ISSN: 1523-7060 cited in the application	1
Y	the whole document	1-5
P,X	EP 1 088 887 A (DSM NV) 4 April 2001 (2001-04-04)	1
P,Y	the whole document	1-5
Y	VAN UNEN, DIRK-JAN ET AL: "Crown ether activation of crosslinked subtilisin Carlsberg crystals in organic solvents" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1: ORGANIC AND BIO-ORGANIC CHEMISTRY (20), 1998, pages 3341-3343, XP002202482 the whole document	1-5
Y	KHALAF ET AL.: "Cross-linked enzyme crystals as highly active catalysts in organic solvents" J. AM. CHEM. SOC., vol. 118, 1996, pages 5494-5495, XP002202483 the whole document	1-5
Y	SHARON C ET AL: "Purification and characterization of a lipase from Pseudomonas aeruginosa KKA-5 and its role in castor oil hydrolysis." JOURNAL OF INDUSTRIAL MICROBIOLOGY & BIOTECHNOLOGY, vol. 20, no. 5, May 1998 (1998-05), pages 304-307, XP001083606 ISSN: 1367-5435 page 306, column 2, paragraph 3; table 2	1-5
Y	DATABASE WPI Week 199803 Derwent Publications Ltd., London, GB; AN 1998-021952 XP002202391 & JP 08 113593 A ((UMEM-I) UMEMOTO M), 7 May 1996 (1996-05-07) abstract	1-5

-/-

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 02/00079

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 17504 A (BASF AG ;BALKENHOHL FRIEDHELM (DE); HAUER BERNHARD (DE); LADNER WO) 29 June 1995 (1995-06-29) the whole document -----	1-5
A	US 4 665 028 A (AMOTZ SHMUEL) 12 May 1987 (1987-05-12) the whole document -----	1-5

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 02/00079

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1088887	A	04-04-2001	EP 1088887 A1	04-04-2001
JP 8113593	A	07-05-1996	JP 8114599 A	07-05-1996
WO 9517504	A	29-06-1995	DE 4344211 A1 WO 9517504 A1	29-06-1995 29-06-1995
US 4665028	A	12-05-1987	BE 897926 A1 CA 1205402 A1 DE 3336257 A1 DK 442883 A ,B, ES 526247 D0 ES 8504247 A1 FI 833615 A ,B, FR 2534272 A1 GB 2129809 A ,B GB 2128620 A ,B IE 56079 B1 IT 1163947 B JP 1723280 C JP 4007192 B JP 59130184 A KR 9007631 B1 NL 8303415 A ,B,	05-04-1984 03-06-1986 12-04-1984 07-04-1984 01-04-1985 01-07-1985 07-04-1984 13-04-1984 23-05-1984 02-05-1984 10-04-1991 08-04-1987 24-12-1992 10-02-1992 26-07-1984 17-10-1990 01-05-1984

THIS PAGE BLANK (USPTO)